






















the 1990s, the incidence of *S. flexneri* has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported as the most common serotype in children with acute bacterial dysentery [11].

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. In the 1970s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [12]. In the 1980s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [13].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [14]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [15].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [16]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [17].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [18]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [19].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [20]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [21].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [22]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [23].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [24]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [25].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [26]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [27].

The first of these is the *Journal of the American Medical Association* (JAMA), which has been the most influential of the medical journals in the United States since its founding in 1883. It is a weekly publication that covers a wide range of medical topics, from clinical medicine to public health. The JAMA is known for its high standards of scientific rigor and its commitment to providing the most up-to-date information to its readers. It is also known for its editorial independence and its refusal to accept payment for the publication of articles.

The second of these journals is the *New England Journal of Medicine* (NEJM), which was founded in 1812 and is the oldest of the three. It is a weekly publication that covers a wide range of medical topics, from clinical medicine to public health. The NEJM is known for its high standards of scientific rigor and its commitment to providing the most up-to-date information to its readers. It is also known for its editorial independence and its refusal to accept payment for the publication of articles.

The third of these journals is the *Lancet*, which was founded in 1823 and is the oldest of the three. It is a weekly publication that covers a wide range of medical topics, from clinical medicine to public health. The Lancet is known for its high standards of scientific rigor and its commitment to providing the most up-to-date information to its readers. It is also known for its editorial independence and its refusal to accept payment for the publication of articles.

These three journals are the most influential of the medical journals in the United States, and they are also the most respected. They are known for their high standards of scientific rigor and their commitment to providing the most up-to-date information to their readers. They are also known for their editorial independence and their refusal to accept payment for the publication of articles.















-----

